CADDISFLIES (TRICHOPTERA) COLLECTED IN THE LAKE DISTRICT (CUMBRIA) IN JULY 2006

EMMA ROSS

Department of Entomology, The Natural History Museum, London, SW7 5BD e.ross@nhm.ac.uk

While on holiday in the Lake District during 14–27th July 2006, I took the opportunity to collect caddisflies at a variety of localities. I stayed at Ennerdale, by the River Ehen, which was a good position for visiting different potential caddisfly localities in the western part of the Lake District. This was the perfect time to visit as the rest of the country was experiencing a heatwave and the temperature was generally 5°C lower there than in the rest of England. Most days were fine and sunny, though the waterfalls were not at their best as there had not been much rain.

Prior to going, Ian Wallace (pers. comm.) informed me that there were good records for some of the well visited and popular sites in the southern Lake District e.g. Lake Windermere, but few records from other localities. D.E. Kimmins was evacuated to Wray Castle on Windermere, along with the Natural History Museum collections he cared for over the Second World War and published a major list and a supplement (Kimmins, 1943, 1944). I really wanted to find *Philopotamus montanus* (Donovan) (Philopotamidae) which is only known from fast flowing streams, a habitat that I had not collected from before.

There are four main types of freshwater habitats in the Lake District where caddisflies can be found: (1) large lakes, which often have the word 'water' as part of their name; (2) ponds, usually at high altitude which are called 'tarns'; (3) rivers, and 4) fast flowing streams, which are called 'becks'. The latter can have waterfalls, which often have the word 'force' as part of the name and may have cut a chasm know as a 'gill' or 'ghyll'. There are two very good guides to the tarns and waterfalls by Blair (2002, 2003), that proved very useful. A guide to Lake District walks by Conduit, Coates and Taylor (2006) was also useful.

Most collecting was done by just tubing specimens as they were seen, though a net was used for sweeping. A total of 30 species from 22 localities was found; only one or two examples of a taxon were collected from each locality. I was able to take photographs of many living specimens, usually the following morning when they were calmer (e.g. Plate 2, Fig. 2, *Adicella reducta* (McLachlan)). Some of the photos will be published by Barnard & Ross (in press).

The localities and species recorded are listed in Table 1 in alphabetical order. For eight of the localities marked with an asterisk, there are no previous records on the database of the national caddisfly recording scheme (Wallace pers. comm.). Localities owned by the National Trust are indicated by (NT).

All of the species are common according to Wallace (1991). These records have been added to the Cumbrian species list on the Virtual Fauna of Lakeland website (http://www.lakelandwildlife.co.uk).

Philopotamus montanus was found at three localities, all at the bases of waterfalls where it is cool and shaded. Although common, this species proved remarkably difficult to catch. They were very active, fast flying and very difficult to follow in dappled shade and over the slippery rocks. When they land on mossy rocks they virtually disappear as their yellow and brown mottled wings make them very well camouflaged. The easiest way to spot them is to get under overhanging vegetation

Table 1. Localities in the Lake District where caddisflies were collected in July 2006. Localities for which there are no previous caddisfly records are indicated by an asterisk.

Aira Beck

Aira Force and below High Force*

[NY 400 207] (NT), 25.vii.06:

Adicella reducta (McLachlan), Beraea maurus (Curtis), Tinodes maclachlani Kimmins

Bassenthwaite Lake

North end [NY 201 319], 21.vii.06:

Tinodes waeneri (L.)

Burnmoor Tarn*

[NY 18 04] (NT), 23.vii.06:

Athripsodes albifrons (L.), Athripsodes cinereus (Curtis), Cyrnus trimaculatus (Curtis), Lype phaeopa (Stephens), Mystacides nigra (L.), Tinodes waeneri

Outflow from Burnmoor Tarn into Whillan Beck [NY 188 044]: Hydropsyche siltalai Döhler

Buttermere

[NY 17 16–18 15] (NT), 24.vii.06: Agrypnia varia (Fabr.), Goera pilosa (Fabr.)

Buttermere Dubs*

A waterway that connects the lakes of Crummock Water and Buttermere [NY 170 164], 24.vii.06: Neureclipsis bimaculata (L.), Oecetis testacea (Curtis), Tinodes waeneri

Cogra Moss

Also known as Arlecdon Reservoir [NY 09 19], 26.vii.06:

Athripsodes aterrimus (Stephens), Cyrnus flavidus McLachlan, Cyrnus trimaculatus, Oecetis lacustris (Pictet)

Dub Beck

[NY 134 211], 27.vii.06: Athripsodes albifrons

Ennerdale Bridge

Stream next to Fox & Hounds pub [NY 069 159], 26.vii.06: Odontocerum albicorne (Scopoli)

Ennerdale Water (NT)

North [NY 103 159] and east sides [NY 124 139], 15.vii.06:

Cyrnus trimaculatus, Polycentropus flavomaculatus (Pictet)

South side, with small shaded streams flowing into it [NY 112 143] (NT), 15.vii.06:

Diplectrona felix McLachlan

West side [NY 089 155], 26.vii.06: Cyrnus trimaculatus

Friar's Crag, Derwent Water [NY 264 222], 21.vii.06: Cyrnus trimaculatus, Polycentropus flavomaculatus

Holme Beck*

At the base of Holme Force [NY 119 214] (NT), 27.vii.06: Drusus annulatus (Stephens)

Loweswater (NT)

North-east side [NY 125 219], 24.vii.06: Athripsodes cinereus

South-west side [NY 125 212], 27.vii.06: Polycentropus flavomaculatus, Tinodes waeneri

Lingcove Beck*

At base of a waterfall [NY 228 037] (NT), 16.vii.06:

Lepidostoma hirtum (Fabr.), Oecetis testacea

Table 1. (Continued)

River Derwent, Cockermouth

[NY 122 309], 20.vii.06: Rhyacophila dorsalis (Curtis)

River Ehen

[NY 060 158], 19.vii.06:

Beraea maurus, Lepidostoma hirtum, Odontocerum albicorne, Oecetis testacea 26 vii 06:

Cyrnus trimaculatus, Glossosoma conformis Neboiss, Odontocerum albicorne, Psychomyia pusilla (Fabr.), Rhyacophila dorsalis (Curtis), Silo pallipes (Fabr.)

Scale Beck*

At the base of Scale Force [NY 152 173] (NT), 24.vii.06: *Philopotamus montanus* (Donovan)

Stanley Gill*

Below Stanley Force [SD 175 998], 16.vii.06: Philopotamus montanus, Sericostoma personatum (Spence)

Sprinkling Tarn*

[NY 228 091] (NT), 18.vii.06: Agrypnia obsoleta (Hagen), Cyrnus flavidus, Mystacides azurea (L.), Polycentropus flavomaculatus

Sty Head bog

[NY 220 094] (NT), 18.vii.06: Agrypnia obsoleta

Ullswater

South end, near Glenridding Pier [NY 390 169] (NT), 25.vii.06: Athripsodes aterrimus, Tinodes waeneri

Watendlath Beck

Base of Lodore Falls [NY 265 188] (NT), 21.vii.06: Diplectrona felix, Lepidostoma hirtum, Philopotamus montanus

and look upwards towards a bright sky. They often sit on the top of the leaves and can be seen as a silhouette through translucent leaves.

I would like to thank Peter Barnard (NHM) for confirmation of identifications, Ian Wallace for information on localities, the National Trust for access, Steve Hewitt (Tullie House Museum) for information about Cumbrian caddisfly records and Andrew Ross (NHM) for general help.

REFERENCES

Barnard, P. & Ross, E. (in press). A guide to the adult caddisflies or sedge flies (Trichoptera). AIDGAP. Field Studies Council.

Blair, D. 2002. Exploring Lakeland Waterfalls. Lakeland Moor Press. 188 pp.

Blair, D. 2003. Exploring Lakeland Tarns. Revised Edition. Lakeland Moor Press. 223 pp.
Conduit, B., Coates, N. & Taylor, H. 2006. Lake District Walks. Revised Edition. Ordnance Survey Pathfinder Guide, No. 13. Jarrold Publishing. 95 pp.

Kimmins, D.E. 1943. A list of the Trichoptera (caddisflies) of the Lake District with distributional and seasonal data. *Journal of the Society for British Entomology* 2: 136–157. Kimmins, D.E. 1944. Supplementary notes on the Trichoptera of the English Lake District.

Entomologist 77: 81–83.

Wallace, I.D. 1991. A review of the Trichoptera of Great Britain. *Joint Nature Conservation Committee Report*, No. 32, 61 pp.